

On Octonary Codes and their Covering Radii

Manoj K. Raut and Manish K. Gupta

Dhirubhai Ambani Institute of Information and Communication Technology,
Gandhinagar, Gujarat-382007

{manoj_raut@daiict.ac.in, mankg@daiict.ac.in}
<http://www.daiict.ac.in>

Abstract. This paper introduces new reduction and torsion codes for an octonary code and determines their basic properties. These could be useful for the classification of self-orthogonal and self dual codes over \mathbb{Z}_8 . We also focus our attention on covering radius problem of octonary codes. In particular, we determine lower and upper bounds of the covering radius of several classes of Repetition codes, Simplex codes of Type α and Type β and their duals, MacDonald codes, and Reed-Muller codes over \mathbb{Z}_8 .

Keywords: Octonary codes, Reduction codes, Torsion codes, Covering radius, codes over rings, Simplex codes, MacDonald codes, Reed-Muller codes.

1 Introduction

Codes over finite rings is an interesting area investigated by many researchers in the last two decades [2, 3, 5–7, 9, 14, 17–19, 22–24, 28–30, 32–34]. In particular, Octonary codes have received attention by many researchers [2, 5, 9, 20, 22, 24, 32]. For any octonary linear code, we introduce binary and quaternary (over \mathbb{Z}_4) reduction and torsion codes and study their basic properties with respect to self-orthogonality and self-duality. One of the other important properties of error correcting codes is covering radius. The covering radius of binary linear codes have been studied in [10, 11]. It is shown in [4, 11] that the problem of computing covering radii of codes is both NP-hard and Co-NP hard. Infact, this problem is strictly harder than any NP-complete problem, unless NP=co-NP. Recently, the covering radius of codes over \mathbb{Z}_4 has been investigated with respect to Lee and Euclidean distances [1]. Several upper and lower bounds on the covering radius of codes has been studied in [1]. More recently covering radius of codes over \mathbb{Z}_{2^s} have been defined in [27] and upper and lower bounds on the covering radius of several classes of codes over \mathbb{Z}_4 have been obtained [27]. We extend some of these results to octonary codes in this paper.

A linear code \mathcal{C} , of length n , over \mathbb{Z}_8 is an additive subgroup of \mathbb{Z}_8^n . An element of \mathcal{C} is called a *codeword* of \mathcal{C} and a *generator matrix* of \mathcal{C} is a matrix whose rows generate \mathcal{C} . The *Hamming weight* $w_H(\mathbf{x})$ of a vector \mathbf{x} in \mathbb{Z}_8^n is the

number of non-zero components of \mathbf{x} . The *Homogeneous weight* $w_{HW}(\mathbf{x})$ [13] of a vector $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{Z}_8^n$ is given by $\sum_{i=1}^n w_{HW}(x_i)$ where

$$w_{HW}(x_i) = \begin{cases} 2, & x_i \neq 4 \\ 4, & x_i = 4. \end{cases} \quad (1)$$

The *Lee weight* $w_L(\mathbf{x})$ of a vector $\mathbf{x} \in \mathbb{Z}_8^n$ is $\sum_{i=1}^n \min\{x_i, 8 - x_i\}$. The *Euclidean weight* $w_E(\mathbf{x})$ of a vector $\mathbf{x} \in \mathbb{Z}_8^n$ is $\sum_{i=1}^n \min\{x_i^2, (8 - x_i)^2\}$.

The Hamming, Homogeneous, Lee and Euclidean distances $d_H(\mathbf{x}, \mathbf{y})$, $d_{HW}(\mathbf{x}, \mathbf{y})$, $d_L(\mathbf{x}, \mathbf{y})$, and $d_E(\mathbf{x}, \mathbf{y})$ between two vectors \mathbf{x} and \mathbf{y} are $w_H(\mathbf{x} - \mathbf{y})$, $w_{HW}(\mathbf{x} - \mathbf{y})$, $w_L(\mathbf{x} - \mathbf{y})$ and $w_E(\mathbf{x} - \mathbf{y})$, respectively. The minimum Hamming, Homogeneous, Lee and Euclidean weights, d_H, d_{HW}, d_L and d_E of \mathcal{C} are the smallest Hamming, Homogeneous, Lee and Euclidean weights among all non-zero codewords of \mathcal{C} respectively. One can define an isometry called the generalized *Gray map* $\phi : \mathbb{Z}_8^n \rightarrow \mathbb{Z}_2^{4n}$ as a coordinate-wise extension of the function from \mathbb{Z}_8 to \mathbb{Z}_2^4 defined by $0 \rightarrow (0, 0, 0, 0), 1 \rightarrow (0, 1, 0, 1), 2 \rightarrow (0, 0, 1, 1), 3 \rightarrow (0, 1, 1, 0), 4 \rightarrow (1, 1, 1, 1), 5 \rightarrow (1, 0, 1, 0), 6 \rightarrow (1, 1, 0, 0), 7 \rightarrow (1, 0, 0, 1)$ [8]. The image $\phi(\mathcal{C})$, of a linear code \mathcal{C} over \mathbb{Z}_8 of length n by the generalized Gray map, is a binary code of length $4n$ [28].

Let $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_n)$ be two vectors in \mathbb{Z}_8^n . Then the inner product of \mathbf{x} and \mathbf{y} is defined by $\mathbf{x} \cdot \mathbf{y} = (x_1 y_1 + x_2 y_2 + \dots + x_n y_n) \pmod{8}$. The *dual code* \mathcal{C}^\perp of \mathcal{C} is defined as $\{\mathbf{x} \in \mathbb{Z}_8^n \mid \mathbf{x} \cdot \mathbf{y} = 0 \text{ for all } \mathbf{y} \in \mathcal{C}\}$ where $\mathbf{x} \cdot \mathbf{y}$ is the inner product of \mathbf{x} and \mathbf{y} . \mathcal{C} is *self-orthogonal* if $\mathcal{C} \subseteq \mathcal{C}^\perp$ and \mathcal{C} is *self-dual* if $\mathcal{C} = \mathcal{C}^\perp$.

Two codes are said to be *equivalent* if one can be obtained from the other by permuting the coordinates and (if necessary) changing the signs of certain coordinates. Codes differing by only a permutation of coordinates are called *permutation-equivalent*. Let $\mathcal{C} \subseteq \mathbb{Z}_8^n$. If \mathcal{C} has M codewords and minimum Homogeneous and Euclidean distances d_{HW} and d_E respectively then \mathcal{C} is called an (n, M, d_{HW}, d_E) code. For more details about octonary codes reader is referred to any of the papers from [20, 24].

This paper is organized as follows. In Section 2, we define new torsion and reduction codes for an octonary code and obtained their basic properties. In Section 3 we present some results of covering radius of octonary codes. Section 4, we discuss about covering radius of Octonary repetition codes. Octonary simplex codes of type α and β is discussed in Section 5. In Section 6, we consider MacDonald codes \mathbb{Z}_8 . Finally Section 7 considers Reed-Muller codes and Section 8 considers Octocode. Last section concludes the paper.

2 Reduction and Torsion Codes

The standard form of generator matrix G of the linear code \mathcal{C} over \mathbb{Z}_8 [20] is of the form

$$G = \begin{pmatrix} I_{k_0} & A_{0,1} & A_{0,2} & A_{0,3} \\ 0 & 2I_{k_1} & 2A_{1,2} & 2A_{1,3} \\ 0 & 0 & 4I_{k_2} & 4A_{2,3} \end{pmatrix}, \quad (2)$$

where the matrices $A_{i,j}$ are binary matrices for $i > 0$. A code with a generator matrix in this form is of type $\{k_0, k_1, k_2\}$ and has $8^{k_0} 4^{k_1} 2^{k_2}$ vectors.

The matrix (2) can also be written in the following form over \mathbb{Z}_8 :

$$G = \begin{pmatrix} I_{k_0} & B_{0,1} + 2B_{0,1}^1 + 4B_{0,1}^2 & B_{0,2} + 2B_{0,2}^1 + 4B_{0,2}^2 & B_{0,3} + 2B_{0,3}^1 + 4B_{0,3}^2 \\ 0 & 2I_{k_1} & 2A_{1,2} & 2A_{1,3} \\ 0 & 0 & 4I_{k_2} & 4A_{2,3} \end{pmatrix}, \quad (3)$$

where $B_{0,i}$, $B_{0,i}^1$, and $B_{0,i}^2$ are binary matrices for $i > 0$ and the matrices $A_{i,j}$ are binary matrices for $i > 0$. For a quaternary linear code one can define reduction code and torsion code [14, 28]. These codes have been generalized for a linear code \mathcal{C} over \mathbb{Z}_8 in the form of four binary torsion/reduction codes in [20]. For $0 \leq i \leq 3$,

$$Tor_i(\mathcal{C}) = \{v \pmod{2} \mid 2^i v \in \mathcal{C}\}. \quad (4)$$

The generator matrices of $Tor_0(\mathcal{C})$, $Tor_1(\mathcal{C})$, $Tor_2(\mathcal{C})$ are the following three binary matrices:

$$G_{Tor_0} = (I_{k_0} \ B_{0,1} + 2B_{0,1}^1 + 4B_{0,1}^2 \ B_{0,2} + 2B_{0,2}^1 + 4B_{0,2}^2 \ B_{0,3} + 2B_{0,3}^1 + 4B_{0,3}^2) \quad (5)$$

$$G_{Tor_1} = \begin{pmatrix} I_{k_0} & B_{0,1} + 2B_{0,1}^1 + 4B_{0,1}^2 & B_{0,2} + 2B_{0,2}^1 + 4B_{0,2}^2 & B_{0,3} + 2B_{0,3}^1 + 4B_{0,3}^2 \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \end{pmatrix} \quad (6)$$

$$G_{Tor_2} = \begin{pmatrix} I_{k_0} & B_{0,1} + 2B_{0,1}^1 + 4B_{0,1}^2 & B_{0,2} + 2B_{0,2}^1 + 4B_{0,2}^2 & B_{0,3} + 2B_{0,3}^1 + 4B_{0,3}^2 \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \\ 0 & 0 & I_{k_2} & A_{2,3} \end{pmatrix} \quad (7)$$

where

$$|\mathcal{C}| = |Tor_0(\mathcal{C})| |Tor_1(\mathcal{C})| |Tor_2(\mathcal{C})| = 2^{3k_0+2k_1+k_2} \quad (8)$$

The reduction and torsion code of quaternary linear code can also be generalized for linear codes over \mathbb{Z}_8 in another interesting way. We define two binary (over \mathbb{Z}_2) torsion codes and two quaternary (\mathbb{Z}_4) torsion codes for a given linear code over \mathbb{Z}_8 as follows:

$$\begin{aligned} \mathcal{C}^{(1)} &= \{c \pmod{2} \mid c \in \mathcal{C}\}, \\ \mathcal{C}^{(2)} &= \{c \pmod{4} \mid c \in \mathcal{C}\}, \\ \mathcal{C}^{(3)} &= \{c \mid 2c \in \mathcal{C}\}, \\ \mathcal{C}^{(4)} &= \{c \mid 4c \in \mathcal{C}\}. \end{aligned} \quad (9)$$

The generator matrices $G^{(1)}, G^{(2)}, G^{(3)}, G^{(4)}$ of $\mathcal{C}^{(1)}, \mathcal{C}^{(2)}, \mathcal{C}^{(3)}, \mathcal{C}^{(4)}$ are obtained from equation (3) as follows:

$$G^{(1)} = (I_{k_0} \ B_{0,1} \ B_{0,2} \ B_{0,3}), \quad (10)$$

$$G^{(2)} = \begin{pmatrix} I_{k_0} & B_{0,1} + 2B_{0,1}^1 & B_{0,2} + 2B_{0,2}^1 & B_{0,3} + 2B_{0,3}^1 \\ 0 & 2I_{k_1} & 2A_{1,2} & 2A_{1,3} \end{pmatrix}, \quad (11)$$

$$G^{(3)} = \begin{pmatrix} I_{k_0} & B_{0,1} + 2B_{0,1}^1 & B_{0,2} + 2B_{0,2}^1 & B_{0,3} + 2B_{0,3}^1 \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \\ 0 & 0 & 2I_{k_2} & 2A_{2,3} \end{pmatrix}, \quad (12)$$

$$G^{(4)} = \begin{pmatrix} I_{k_0} & B_{0,1} & B_{0,2} & B_{0,3} \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \\ 0 & 0 & I_{k_2} & A_{2,3} \end{pmatrix}. \quad (13)$$

Note that the number of elements of \mathcal{C} is $8^{k_0} 4^{k_1} 2^{k_2} = 2^{3k_0+2k_1+k_2}$. The number of elements of $\mathcal{C}^{(1)}, \mathcal{C}^{(2)}, \mathcal{C}^{(3)}, \mathcal{C}^{(4)}$ are $2^{k_0}, 4^{k_0} 2^{k_1}, 4^{k_0+k_1} 2^{k_2}$ and $2^{k_0+k_1+k_2}$ respectively. Thus for a linear code \mathcal{C} over \mathbb{Z}_8 we have the following relationship.

Proposition 1.

$$|\mathcal{C}| = |\mathcal{C}^{(1)}| \times |\mathcal{C}^{(3)}| = |\mathcal{C}^{(2)}| \times |\mathcal{C}^{(4)}|.$$

Note that if $k_1 = k_2 = 0$ then $\mathcal{C}^{(2)} = \mathcal{C}^{(3)}$. It is easy to observe the following.

Proposition 2.

$$\mathcal{C}^{(1)} \subseteq \mathcal{C}^{(4)} \text{ and } \mathcal{C}^{(2)} \subseteq \mathcal{C}^{(3)}.$$

Next result is a simple generalization of self-orthogonality characterization from [25]. For $\mathbf{c} \in \mathcal{C}$ and $0 \leq i \leq 7$, let $w_i(\mathbf{c})$ denotes the composition of symbol i in the codeword \mathbf{c} .

Proposition 3. *A linear code \mathcal{C} over \mathbb{Z}_8 is self-orthogonal iff each generator matrix of \mathcal{C} has all its rows $\omega_1 + \omega_3 + \omega_5 + \omega_7 + 4\omega_2 + 4\omega_6 = 0 \pmod{8}$ and every pair of rows of the generator matrix is orthogonal.*

Proof. Proof is straightforward. \square

Now we determine few relationships among various reduction and torsion codes if code \mathcal{C} is self-orthogonal or self-dual.

Proposition 4. *If \mathcal{C} is a self-orthogonal code over \mathbb{Z}_8 then $\mathcal{C}^{(1)}, \mathcal{C}^{(4)}$ are self-orthogonal codes over \mathbb{Z}_2 and $\mathcal{C}^{(2)}, \mathcal{C}^{(3)}$ are self-orthogonal codes over \mathbb{Z}_4 .*

Proof. The self-orthogonality of $\mathcal{C}^{(1)}, \mathcal{C}^{(2)}$ follows from [20]. It remains to see the self-orthogonality of $\mathcal{C}^{(3)}$, and $\mathcal{C}^{(4)}$. Let $v \in \mathcal{C}^{(3)}$. By definition of $\mathcal{C}^{(3)}$ we have $2v \in \mathcal{C}$. As \mathcal{C} is self orthogonal, $\langle 2v, u \rangle = 0 \pmod{4}$ for all $u \in \mathcal{C}$. So $2 \sum v_i u_i \equiv 0 \pmod{8}$ for all $u \in \mathcal{C}$. Then $\sum v_i u_i \equiv 0 \pmod{4}$ for all $u \in \mathcal{C}$. This implies $\langle v, u \rangle = 0$ for all $u \in \mathcal{C}^{(3)}$ as $\mathcal{C}^{(3)}$ is a code over \mathbb{Z}_4 . So $v \in \mathcal{C}^{(3)\perp}$. Hence $\mathcal{C}^{(3)}$ is self orthogonal. The self-orthogonality of $\mathcal{C}^{(4)}$ can be proved similarly. \square

Proposition 5. *If \mathcal{C} is self orthogonal code over \mathbb{Z}_8 then $\mathcal{C}^{(4)} \subseteq \mathcal{C}^{(1)\perp}$ and $\mathcal{C}^{(3)} \subseteq \mathcal{C}^{(2)\perp}$.*

Proof. Let $v \in \mathcal{C}^{(4)}$. By definition of $\mathcal{C}^{(4)}$, we have $4v \in \mathcal{C}$. As \mathcal{C} is self-orthogonal, $\langle 4v, u \rangle = 0 \pmod{8}$ for all $u \in \mathcal{C}$. So $4 \sum v_i u_i = 0 \pmod{8}$ for all $u \in \mathcal{C}$. $\sum v_i u_i = 0 \pmod{2}$ for all $u \in \mathcal{C}$. $\sum (v_i \pmod{2})(u_i \pmod{2}) = 0 \pmod{2}$ for all $u \pmod{2} \in \mathcal{C}$. As $\mathcal{C}^{(4)}$ is a code over \mathbb{Z}_2 , $v_i \pmod{2} = v_i$ and as $\mathcal{C}^{(1)}$ is a code over \mathbb{Z}_2 , $\sum v_i (u_i \pmod{2}) = 0 \pmod{2}$ for all $u \in \mathcal{C}^{(1)}$. So $\langle v, u \rangle = 0 \pmod{2}$ for all $u \in \mathcal{C}^{(1)}$. This implies $v \in \mathcal{C}^{(1)\perp}$. The second inclusion $\mathcal{C}^{(3)} \subseteq \mathcal{C}^{(2)\perp}$ can be proved similarly. \square

Proposition 6. *If \mathcal{C} is self-dual over \mathbb{Z}_8 then $\mathcal{C}^{(4)} = \mathcal{C}^{(1)\perp}$ and $\mathcal{C}^{(3)} = \mathcal{C}^{(2)\perp}$.*

Proof. We know that $\mathcal{C}^{(4)} \subseteq \mathcal{C}^{(1)\perp}$ from Proposition 5. It remains to show $\mathcal{C}^{(1)\perp} \subseteq \mathcal{C}^{(4)}$. Let $v \in \mathcal{C}^{(1)\perp}$. So $\langle v, w \rangle \equiv 0 \pmod{2}$ for all $w \in \mathcal{C}^{(1)}$. $\sum v_i w_i \equiv 0 \pmod{2}$ for all $w \in \mathcal{C}^{(1)}$. $4 \sum v_i w_i \equiv 0 \pmod{8}$ for all $w \in \mathcal{C}$. $\sum 4v_i w_i \equiv 0 \pmod{8}$ for all $w \in \mathcal{C}$. $\langle 4v, w \rangle = 0 \pmod{8}$ for all $w \in \mathcal{C}$. $4v \in \mathcal{C}^\perp = \mathcal{C}$. This implies $v \in \mathcal{C}^{(4)}$. Hence proved. Proof of second result is similar. \square

We know that $\mathcal{C}^{(2)}$ and $\mathcal{C}^{(3)}$ are codes over \mathbb{Z}_4 . Thus it is natural to consider the torsion and reduction code of $\mathcal{C}^{(2)}$ and $\mathcal{C}^{(3)}$. We get the following:

$$\begin{aligned} \mathcal{C}^{(21)} &= \{c \pmod{2} \mid c \in \mathcal{C}^{(2)}\}, \\ \mathcal{C}^{(22)} &= \{c \mid 2c \in \mathcal{C}^{(2)}\}, \\ \mathcal{C}^{(31)} &= \{c \pmod{2} \mid c \in \mathcal{C}^{(3)}\}, \\ \mathcal{C}^{(32)} &= \{c \mid 2c \in \mathcal{C}^{(3)}\}. \end{aligned} \tag{14}$$

The generator matrices $G^{(21)}, G^{(22)}, G^{(31)}, G^{(32)}$ of $\mathcal{C}^{(21)}, \mathcal{C}^{(22)}, \mathcal{C}^{(31)}, \mathcal{C}^{(32)}$ are obtained from (11) and (12) as follows:

$$G^{(21)} = (I_{k_0} \ B_{0,1} \ B_{0,2} \ B_{0,3}) = G^{(1)}, \tag{15}$$

$$G^{(22)} = \begin{pmatrix} I_{k_0} & B_{0,1} & B_{0,2} & B_{0,3} \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \end{pmatrix}, \tag{16}$$

$$G^{(31)} = \begin{pmatrix} I_{k_0} & B_{0,1} & B_{0,2} & B_{0,3} \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \end{pmatrix} = G^{(22)}, \tag{17}$$

$$G^{(32)} = \begin{pmatrix} I_{k_0} & B_{0,1} & B_{0,2} & B_{0,3} \\ 0 & I_{k_1} & A_{1,2} & A_{1,3} \\ 0 & 0 & I_{k_2} & A_{2,3} \end{pmatrix}. \tag{18}$$

All the codes $\mathcal{C}^{(21)}, \mathcal{C}^{(22)}, \mathcal{C}^{(31)}, \mathcal{C}^{(32)}$ are codes over \mathbb{Z}_2 . It is easy to see the following results from their generator matrices:

Proposition 7. *If \mathcal{C} is a code over \mathbb{Z}_8 then $\mathcal{C}^{(21)} = \mathcal{C}^{(1)}, \mathcal{C}^{(1)} \subseteq \mathcal{C}^{(22)}, \mathcal{C}^{(31)} = \mathcal{C}^{(22)}, \mathcal{C}^{(31)} \subseteq \mathcal{C}^{(32)}$, and $\mathcal{C}^{(32)} = \mathcal{C}^{(4)}$.*

Also it is natural to obtain the following from [14].

Proposition 8. *If \mathcal{C} is self-orthogonal over \mathbb{Z}_8 then*

1. $\mathcal{C}^{(21)} \subseteq \mathcal{C}^{(22)} \subseteq \mathcal{C}^{(21)\perp}$.
2. $\mathcal{C}^{(31)} \subseteq \mathcal{C}^{(32)} \subseteq \mathcal{C}^{(31)\perp}$.

Thus we have interesting family of codes from a linear octonary codes having beautiful inclusions.

3 Covering Radius of Octonary Codes

In this section, first we collect some known facts of the covering radius of codes over \mathbb{Z}_8 with respect to Homogeneous and Euclidean distances [27] and then derive some of its properties. Let d be either a Homogeneous distance or Euclidean distance. Then the covering radius of code \mathcal{C} over \mathbb{Z}_8 with respect to distance d is given by

$$r_d(\mathcal{C}) = \max_{\mathbf{u} \in \mathbb{Z}_8^n} \left\{ \min_{\mathbf{c} \in \mathcal{C}} d(\mathbf{u}, \mathbf{c}) \right\}.$$

We can easily see [27] that $r_d(\mathcal{C})$ is the minimum value r_d such that

$$\mathbb{Z}_8^n = \cup_{\mathbf{c} \in \mathcal{C}} S_{r_d}(\mathbf{c})$$

where

$$S_{r_d}(\mathbf{u}) = \{\mathbf{v} \in \mathbb{Z}_8^n \mid d(\mathbf{u}, \mathbf{v}) \leq r_d\}$$

for any element $\mathbf{u} \in \mathbb{Z}_8^n$.

The coset of \mathcal{C} is the translate $\mathbf{u} + \mathcal{C} = \{\mathbf{u} + \mathbf{c} \mid \mathbf{c} \in \mathcal{C}\}$ where $\mathbf{u} \in \mathbb{Z}_8^n$. A vector of least weight in a coset is called a *coset leader*. The following proposition is well known [27].

Proposition 9. *The covering radius of \mathcal{C} with respect to the general distance d is the largest minimum weight among all cosets.*

Proposition 10. *For any octonary code over \mathbb{Z}_8 ,*

$$\begin{aligned} \frac{1}{2}r_{HW}(\mathcal{C}) &\leq r_E(\mathcal{C}) \leq 5r_{HW}(\mathcal{C}), \\ r_L(\mathcal{C}) &\leq r_E(\mathcal{C}), \\ r_{HW}(\mathcal{C}) &\leq 2r_L(\mathcal{C}). \end{aligned} \tag{19}$$

Proof. We observe that $\frac{1}{2}d_{HW}(\mathbf{x}, \mathbf{y}) \leq d_E(\mathbf{x}, \mathbf{y}) \leq 5d_{HW}(\mathbf{x}, \mathbf{y})$, so the first inequality follows. As $d_L(\mathbf{x}, \mathbf{y}) \leq d_E(\mathbf{x}, \mathbf{y})$, the second inequality follows. Further the third inequality follows since we have $d_{HW}(\mathbf{x}, \mathbf{y}) \leq 2d_L(\mathbf{x}, \mathbf{y})$. \square

The following proposition is also well known [27].

Proposition 11. *Let \mathcal{C} be a code over \mathbb{Z}_8 and $\phi(\mathcal{C})$ the generalized Gray map image of \mathcal{C} . Then $r_{HW}(\mathcal{C}) = r_H(\phi(\mathcal{C}))$.*

The following two results are two upper bounds of the covering radius of codes over \mathbb{Z}_8 with respect to Homogeneous weight.

Proposition 12. (Sphere-Covering Bound) *For any code \mathcal{C} of length n over \mathbb{Z}_8 ,*

$$\begin{aligned} \frac{2^{4n}}{|\mathcal{C}|} &\leq \sum_{i=0}^{r_{HW}(\mathcal{C})} \binom{4n}{i}, \\ \frac{2^{4n}}{|\mathcal{C}|} &\leq \sum_{i=0}^{r_E(\mathcal{C})} V_i, \end{aligned} \quad (20)$$

where $\sum_{i=0}^{16n} V_i x^i = (1 + 2x + 2x^4 + 2x^9 + x^{16})^n$.

Proof. The proof of both inequality over \mathbb{Z}_8 is similar to the proof over \mathbb{Z}_4 given in [1] (see also [27]) and hence omitted. \square

Let \mathcal{C} be a code over \mathbb{Z}_8 and let $s(\mathcal{C}^\perp) = |\{i \mid A_i(\mathcal{C}^\perp) \neq 0, i \neq 0\}|$, where $A_i(\mathcal{C}^\perp)$ is the number of codewords of homogenous weight i in \mathcal{C}^\perp .

Theorem 1. (Delsarte Bound) *Let \mathcal{C} be a code over \mathbb{Z}_8 then $r_{HW}(\mathcal{C}) \leq s(\mathcal{C}^\perp)$ and $r_E(\mathcal{C}) \leq 5s(\mathcal{C}^\perp)$.*

Proof. First result is obtained in [27]. Second result follows from [1] and Proposition 10. \square

The following result of Mattson [10] is useful for computing covering radii of codes over rings [27].

Proposition 13. (Mattson) *If \mathcal{C}_0 and \mathcal{C}_1 are codes over \mathbb{Z}_8 generated by matrices G_0 and G_1 respectively and if \mathcal{C} is the code generated by*

$$G = \left(\begin{array}{c|c} 0 & G_1 \\ \hline G_0 & A \end{array} \right),$$

then $r_d(\mathcal{C}) \leq r_d(\mathcal{C}_0) + r_d(\mathcal{C}_1)$ and the covering radius of \mathcal{D} (concatenation of \mathcal{C}_0 and \mathcal{C}_1) satisfies the following

$$r_d(\mathcal{D}) \geq r_d(\mathcal{C}_0) + r_d(\mathcal{C}_1),$$

for all distances d over \mathbb{Z}_8 .

Now we determine a bound on the covering radius of octonary code and its corresponding reduction and torsion codes. The following result is a generalization of Theorem 4.4 of [1].

Theorem 2. *For a code over \mathbb{Z}_8 , let d_1, d_2, d_3, d_4 denote the minimum Hamming distances of linear codes $\mathcal{C}^{(1)}, \mathcal{C}^{(2)}, \mathcal{C}^{(3)}, \mathcal{C}^{(4)}$ respectively. If $d_1 \geq 8, d_2 \geq 18, d_3 \geq \frac{25}{4}, d_4 \geq \frac{25}{16}$ then*

$$\begin{aligned} r_E(\mathcal{C}) &\geq 9 \min \left\{ \left\lfloor \frac{d_1}{8} \right\rfloor, \left\lfloor \frac{d_2}{18} \right\rfloor, 4 \left\lfloor \frac{d_3}{25} \right\rfloor, 16 \left\lfloor \frac{d_4}{25} \right\rfloor \right\}, \\ r_{HW}(\mathcal{C}) &\geq 2 \min \left\{ \left\lfloor \frac{d_1}{8} \right\rfloor, \left\lfloor \frac{d_2}{18} \right\rfloor, 4 \left\lfloor \frac{d_3}{25} \right\rfloor, 16 \left\lfloor \frac{d_4}{25} \right\rfloor \right\}. \end{aligned}$$

Proof. Let $t = \min \left\{ \left\lfloor \frac{d_1}{8} \right\rfloor, \left\lfloor \frac{d_2}{18} \right\rfloor, 4 \left\lfloor \frac{d_3}{25} \right\rfloor, 16 \left\lfloor \frac{d_4}{25} \right\rfloor \right\}$. Hence $t > 0$. Let $\mathbf{x} = (00 \dots 0 \underbrace{44 \dots 4}_t)$. Let \mathcal{C} be a code over \mathbb{Z}_8 . Let $\mathbf{c} = (c_1, c_2, \dots, c_n) \in \mathcal{C}$ such that $c_i = 0$ or 4 . Hence $\frac{\mathbf{c}}{4} \in \mathcal{C}^{(4)}$. So $wt(\mathcal{C}^{(4)}) \geq d_4$ as the minimum Hamming distance of $\mathcal{C}^{(4)}$ is d_4 . Thus $wt(\frac{\mathbf{c}}{4}) \geq d_4 \geq t$. Let $\mathbf{c} = (00 \dots 0 \underbrace{44 \dots 4}_{\geq d_4})$. Hence

$$\begin{aligned} d_E(\mathbf{c}, \mathbf{x}) &= 16(d_4 - t) \geq 9t, \\ d_{HW}(\mathbf{c}, \mathbf{x}) &= 4(d_4 - t) \geq 2t. \end{aligned}$$

Similarly for $\mathbf{c} \in \mathcal{C}$ such that $\frac{\mathbf{c}}{2} \in \mathcal{C}^{(3)}$ we get

$$\begin{aligned} d_E(\mathbf{c}, \mathbf{x}) &\geq 4d_3 - 16t \geq 9t, \\ d_{HW}(\mathbf{c}, \mathbf{x}) &\geq 2d_3 - 4t \geq 2t. \end{aligned}$$

For $\mathbf{c} \in \mathcal{C}$ such that $\mathbf{c} \pmod{4} \in \mathcal{C}^{(2)}$ we have

$$\begin{aligned} d_E(\mathbf{c}, \mathbf{x}) &\geq d_2 - 9t \geq 9t, \\ d_{HW}(\mathbf{c}, \mathbf{x}) &= 2d_2 \geq 2t. \end{aligned}$$

Finally for $\mathbf{c} \in \mathcal{C}$ such that $\mathbf{c} \pmod{2} \in \mathcal{C}^{(1)}$ we have

$$\begin{aligned} d_E(\mathbf{c}, \mathbf{x}) &= d_1 + 8t \geq 9t, \\ d_{HW}(\mathbf{c}, \mathbf{x}) &= 2d_1 \geq 2t. \end{aligned}$$

Hence the result follows. \square

4 Octonary Repetition Codes

A q -ary repetition code \mathcal{C} over a finite field $\mathbb{F}_q = \{\alpha_0 = 0, \alpha_1 = 1, \alpha_2, \alpha_3, \dots, \alpha_{q-2}\}$ is an $[n, 1, n]$ -code $\mathcal{C} = \{\overline{\alpha} \mid \alpha \in \mathbb{F}_q\}$, where $\overline{\alpha} = \{\alpha, \alpha, \dots, \alpha\}$. The covering radius of \mathcal{C} is $\lceil \frac{n(q-1)}{q} \rceil$ [21]. In [27], several classes of repetition codes over \mathbb{Z}_4 have been studied and their covering radius has been obtained. Now we generalize those results for codes over \mathbb{Z}_8 . Consider the repetition codes over \mathbb{Z}_8 . One can define seven basic repetition codes \mathcal{C}_{α_i} , ($1 \leq i \leq 7$) of length n over \mathbb{Z}_8 generated by $G_{\alpha_1} = [\underbrace{11 \dots 1}_n]$, $G_{\alpha_2} = [\underbrace{22 \dots 2}_n]$, $G_{\alpha_3} = [\underbrace{33 \dots 3}_n]$, $G_{\alpha_4} = [\underbrace{44 \dots 4}_n]$, $G_{\alpha_5} = [\underbrace{55 \dots 5}_n]$, $G_{\alpha_6} = [\underbrace{66 \dots 6}_n]$, $G_{\alpha_7} = [\underbrace{77 \dots 7}_n]$. So the repetition codes are $\mathcal{C}_{\alpha_1} = \mathcal{C}_{\alpha_3} = \mathcal{C}_{\alpha_5} = \mathcal{C}_{\alpha_7} = \{(00 \dots 0), (11 \dots 1), (22 \dots 2), (33 \dots 3), (44 \dots 4), (55 \dots 5), (66 \dots 6), (77 \dots 7)\}$, $\mathcal{C}_{\alpha_2} = \mathcal{C}_{\alpha_6} = \{(00 \dots 0), (22 \dots 2), (44 \dots 4), (66 \dots 6)\}$ and $\mathcal{C}_{\alpha_4} = \{(00 \dots 0), (44 \dots 4)\}$. The following theorems determine the covering radius of \mathcal{C}_{α_i} for $1 \leq i \leq 7$.

Theorem 3. $r_E(\mathcal{C}_{\alpha_1}) = r_E(\mathcal{C}_{\alpha_3}) = r_E(\mathcal{C}_{\alpha_5}) = r_E(\mathcal{C}_{\alpha_7}) = \frac{11n}{2}$ and $r_{HW}(\mathcal{C}_{\alpha_1}) = r_{HW}(\mathcal{C}_{\alpha_3}) = r_{HW}(\mathcal{C}_{\alpha_5}) = r_{HW}(\mathcal{C}_{\alpha_7}) = 2n$

Proof. We know that $r_E(\mathcal{C}_{\alpha_i}) = \max_{x \in \mathbb{Z}_8^n} \{d_E(x, \mathcal{C}_{\alpha_i})\}$. Let $\mathbf{x} \in \mathbb{Z}_8^n$. If \mathbf{x} has composition $(\omega_0, \omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6, \omega_7)$ where $\sum_{i=0}^7 \omega_i = n$, then $d_E(\mathbf{x}, \bar{0}) = n - \omega_0 + 3\omega_2 + 8\omega_3 + 15\omega_4 + 8\omega_5 + 3\omega_6$, $d_E(\mathbf{x}, \bar{1}) = n - \omega_1 + 3\omega_3 + 8\omega_4 + 15\omega_5 + 8\omega_6 + 3\omega_7$, $d_E(\mathbf{x}, \bar{2}) = n - \omega_2 + 3\omega_0 + 3\omega_4 + 8\omega_5 + 15\omega_6 + 8\omega_7$, $d_E(\mathbf{x}, \bar{3}) = n - \omega_3 + 8\omega_0 + 3\omega_1 + 3\omega_5 + 8\omega_6 + 15\omega_7$, $d_E(\mathbf{x}, \bar{4}) = n - \omega_4 + 15\omega_0 + 8\omega_1 + 3\omega_2 + 3\omega_6 + 8\omega_7$, $d_E(\mathbf{x}, \bar{5}) = n - \omega_5 + 8\omega_0 + 15\omega_1 + 8\omega_2 + 3\omega_3 + 3\omega_7$, $d_E(\mathbf{x}, \bar{6}) = n - \omega_6 + 3\omega_0 + 8\omega_1 + 15\omega_2 + 8\omega_3 + 3\omega_4$, and $d_E(\mathbf{x}, \bar{7}) = n - \omega_7 + 3\omega_1 + 8\omega_2 + 15\omega_3 + 8\omega_4 + 3\omega_5$. Then

$$d_E(\mathbf{x}, \mathcal{C}_{\alpha_1}) \leq \frac{8n+36(\omega_0+\omega_1+\omega_2+\omega_3+\omega_4+\omega_5+\omega_6+\omega_7)}{8} = \frac{11n}{2}.$$

Thus $r_E(\mathcal{C}_{\alpha_1}) \leq \frac{11n}{2}$.

Let $\mathbf{x} = \underbrace{00\dots 0}_{t_1} \underbrace{11\dots 1}_{t_2} \underbrace{22\dots 2}_{t_3} \underbrace{33\dots 3}_{t_4} \underbrace{44\dots 4}_{t_5} \underbrace{55\dots 5}_{t_6} \underbrace{66\dots 6}_{t_7} \underbrace{77\dots 7}_{n-7t}$ where $t = \lfloor \frac{n}{8} \rfloor$. Then $d_E(\mathbf{x}, \bar{0}) = n + 36t$, $d_E(\mathbf{x}, \bar{1}) = 4n + 12t$, $d_E(\mathbf{x}, \bar{2}) = 9n - 28t$, $d_E(\mathbf{x}, \bar{3}) = 16n - 84t$, $d_E(\mathbf{x}, \bar{4}) = 9n - 28t$, $d_E(\mathbf{x}, \bar{5}) = 4n + 12t$, $d_E(\mathbf{x}, \bar{6}) = n + 36t$, $d_E(\mathbf{x}, \bar{7}) = 44t$. Thus

$$r_E(\mathcal{C}_{\alpha_1}) \geq \frac{44n+36t+12t-28t-84t-28t+12t+36t+44t}{8} = \frac{11n}{2}.$$

Thus $r_E(\mathcal{C}_{\alpha_1}) = r_E(\mathcal{C}_{\alpha_3}) = r_E(\mathcal{C}_{\alpha_5}) = r_E(\mathcal{C}_{\alpha_7}) = \frac{11n}{2}$. The gray map $\phi(\mathcal{C}_{\alpha_1})$ will be a binary repetition code of length $4n$. Thus $r_{HW}(\mathcal{C}_{\alpha_1}) = \lceil \frac{4n(2-1)}{2} \rceil = 2n = r_{HW}(\mathcal{C}_{\alpha_3}) = r_{HW}(\mathcal{C}_{\alpha_5}) = r_{HW}(\mathcal{C}_{\alpha_7})$. \square

Theorem 4. $r_E(\mathcal{C}_{\alpha_2}) = r_E(\mathcal{C}_{\alpha_6}) = 6n$ and $r_{HW}(\mathcal{C}_{\alpha_2}) = r_{HW}(\mathcal{C}_{\alpha_6}) = 2n$

Proof. The proof is similar to proof of Theorem 3, hence omitted.

Theorem 5. $r_E(\mathcal{C}_{\alpha_4}) = 8n$ and $r_{HW}(\mathcal{C}_{\alpha_4}) = 2n$

Proof. The proof is similar to proof of Theorem 3, hence omitted.

In order to determine the covering radius of Simplex code S_k^α over \mathbb{Z}_8 , we have to define a block repetition code over \mathbb{Z}_8 and find its covering radius. Thus the covering radius of the block repetition code $BRep^{m_1+m_2+\dots+m_7}$ ($n = m_1 + m_2 + \dots + m_7$, $M = 8$, $d_{HW} = \min\{2m_1 + 2m_2 + 2m_3 + 4m_4 + 2m_5 + 2m_6 + 2m_7, 2m_1 + 4m_2 + 2m_3 + 2m_5 + 4m_6 + 2m_7, 4m_1 + 4m_3 + 4m_5 + 4m_7\}$, $d_E = \min\{m_1 + 4m_2 + 9m_3 + 16m_4 + 9m_5 + 4m_6 + m_7, 4m_1 + 16m_2 + 4m_3 + 4m_5 + 16m_6 + 4m_7, 9m_1 + 4m_2 + m_3 + 16m_4 + m_5 + 4m_6 + 9m_7, 16m_1 + 16m_3 + 16m_5 + 16m_7\}$) generated by $G = \underbrace{[11\dots 1]_{m_1}}_{m_1} \underbrace{[22\dots 2]_{m_2}}_{m_2} \underbrace{[33\dots 3]_{m_3}}_{m_3} \underbrace{[44\dots 4]_{m_4}}_{m_4} \underbrace{[55\dots 5]_{m_5}}_{m_5} \underbrace{[66\dots 6]_{m_6}}_{m_6} \underbrace{[77\dots 7]_{m_7}}_{m_7}$ is given in the following theorems.

Theorem 6. $r_E(BRep^{m_1+m_2+\dots+m_7}) = \frac{11}{2}(m_1 + m_3 + m_5 + m_7) + 6(m_2 + m_6) + 8m_4$.

Proof. By proposition 13 and Theorem 3, 4, 5 we have $r_E(BRep^{m_1+m_2+\dots+m_7}) \geq \frac{11}{2}(m_1 + m_3 + m_5 + m_7) + 6(m_2 + m_6) + 8m_4$.

On the other hand, let $\mathbf{x} = (\mathbf{x}_1 \mid \mathbf{x}_2 \mid \mathbf{x}_3 \mid \mathbf{x}_4 \mid \mathbf{x}_5 \mid \mathbf{x}_6 \mid \mathbf{x}_7) \in \mathbb{Z}_8^{m_1+m_2+\dots+m_7}$ with $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4, \mathbf{x}_5, \mathbf{x}_6, \mathbf{x}_7$ have compositions (p_0, p_1, \dots, p_7) , (q_0, q_1, \dots, q_7) , (r_0, r_1, \dots, r_7) , (s_0, s_1, \dots, s_7) , (t_0, t_1, \dots, t_7) , (u_0, u_1, \dots, u_7) , (w_0, w_1, \dots, w_7) such that $p_0 + p_1 + \dots + p_7 = m_1$, $q_0 + q_1 + \dots + q_7 = m_2$, $r_0 + r_1 + \dots + r_7 = m_3$, $s_0 + s_1 + \dots + s_7 = m_4$, $t_0 + t_1 + \dots + t_7 = m_5$, $u_0 + u_1 + \dots + u_7 = m_6$, $w_0 + w_1 + \dots + w_7 = m_7$.

$d_E(\mathbf{x}, \bar{0}) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_0 + 3p_2 + 8p_3 + 15p_4 + 8p_5 + 3p_6 - q_0 + 3q_2 + 8q_3 + 15q_4 + 8q_5 + 3q_6 - r_0 + 3r_2 + 8r_3 + 15r_4 + 8r_5 + 3r_6 - s_0 + 3s_2 + 8s_3 + 15s_4 + 8s_5 + 3s_6 - t_0 + 3t_2 + 8t_3 + 15t_4 + 8t_5 + 3t_6 - u_0 + 3u_2 + 8u_3 + 15u_4 + 8u_5 + 3u_6 - w_0 + 3w_2 + 8w_3 + 15w_4 + 8w_5 + 3w_6$, where $\bar{0}$ is the first vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_1) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_1 + 3p_3 + 8p_4 + 15p_5 + 8p_6 + 3p_7 - q_2 + 3q_0 + 3q_4 + 8q_5 + 15q_6 + 8q_7 - r_3 + 8r_0 + 3r_1 + 3r_5 + 8r_6 + 15r_7 - s_4 + 15s_0 + 8s_1 + 3s_2 + 3s_6 + 8s_7 - t_5 + 8t_0 + 15t_1 + 8t_2 + 3t_3 + 3t_7 - u_6 + 3u_0 + 8u_1 + 15u_2 + 8u_3 + 3u_4 - w_7 + 3w_1 + 8w_2 + 15w_3 + 8w_4 + 3w_5$, where $\mathbf{c}_1 = (\underbrace{11\dots 1}_{m_1} \underbrace{22\dots 2}_{m_2} \underbrace{33\dots 3}_{m_3} \underbrace{44\dots 4}_{m_4} \underbrace{55\dots 5}_{m_5} \underbrace{66\dots 6}_{m_6} \underbrace{77\dots 7}_{m_7})$ is the second vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_2) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_2 + 3p_0 + 3p_4 + 8p_5 + 15p_6 + 8p_7 - q_4 + 15q_0 + 8q_1 + 3q_2 + 3q_6 + 8q_7 - r_6 + 3r_0 + 8r_1 + 15r_2 + 8r_3 + 3r_4 - s_0 + 3s_2 + 8s_3 + 15s_4 + 8s_5 + 3s_6 - t_2 + 3t_0 + 3t_4 + 8t_5 + 15t_6 + 8t_7 - u_4 + 15u_0 + 8u_1 + 3u_2 + 3u_6 + 8u_7 - w_6 + 3w_0 + 8w_1 + 15w_2 + 8w_3 + 3w_4$, where $\mathbf{c}_2 = (\underbrace{22\dots 2}_{m_1} \underbrace{44\dots 4}_{m_2} \underbrace{66\dots 6}_{m_3} \underbrace{00\dots 0}_{m_4} \underbrace{22\dots 2}_{m_5} \underbrace{44\dots 4}_{m_6} \underbrace{66\dots 6}_{m_7})$ is the third vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_3) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_3 + 8p_0 + 3p_1 + 3p_5 + 8p_6 + 15p_7 - q_6 + 3q_0 + 8q_1 + 15q_2 + 8q_3 + 3q_4 - r_1 + 3r_3 + 8r_4 + 15r_5 + 8r_6 + 3r_7 - s_4 + 15s_0 + 8s_1 + 3s_2 + 3s_6 + 8s_7 - t_7 + 3t_1 + 8t_2 + 15t_3 + 8t_4 + 3t_5 - u_2 + 3u_0 + 3u_4 + 8u_5 + 15u_6 + 8u_7 - w_5 + 8w_0 + 15w_1 + 8w_2 + 3w_3 + 3w_7$, where $\mathbf{c}_3 = (\underbrace{33\dots 3}_{m_1} \underbrace{66\dots 6}_{m_2} \underbrace{11\dots 1}_{m_3} \underbrace{44\dots 4}_{m_4} \underbrace{77\dots 7}_{m_5} \underbrace{22\dots 2}_{m_6} \underbrace{55\dots 5}_{m_7})$ is the fourth vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_4) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_4 + 15p_0 + 8p_1 + 3p_2 + 3p_6 + 8p_7 - q_0 + 3q_2 + 8q_3 + 15q_4 + 8q_5 + 3q_6 - r_4 + 15r_0 + 8r_1 + 3r_2 + 3r_6 + 8r_7 - s_0 + 3s_2 + 8s_3 + 15s_4 + 8s_5 + 3s_6 - t_4 + 15t_0 + 8t_1 + 3t_2 + 3t_6 + 8t_7 - u_0 + 3u_2 + 8u_3 + 15u_4 + 8u_5 + 3u_6 - w_4 + 15w_0 + 8w_1 + 3w_2 + 3w_6 + 8w_7$, where $\mathbf{c}_4 = (\underbrace{44\dots 4}_{m_1} \underbrace{00\dots 0}_{m_2} \underbrace{44\dots 4}_{m_3} \underbrace{00\dots 0}_{m_4} \underbrace{44\dots 4}_{m_5} \underbrace{00\dots 0}_{m_6} \underbrace{44\dots 4}_{m_7})$ is the fifth vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_5) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_5 + 8p_0 + 15p_1 + 8p_2 + 3p_3 + 3p_7 - q_2 + 3q_0 + 3q_4 + 8q_5 + 15q_6 + 8q_7 - r_7 + 3r_1 + 8r_2 + 15r_3 + 8r_4 + 3r_5 - s_4 + 15s_0 + 8s_1 + 3s_2 + 3s_6 + 8s_7 - t_1 + 3t_3 + 8t_4 + 15t_5 + 8t_6 + 3t_7 - u_6 + 3u_0 + 8u_1 + 15u_2 + 8u_3 + 3u_4 - w_3 + 8w_0 + 3w_1 + 3w_5 + 8w_6 + 15w_7$, where

$\mathbf{c}_5 = (\underbrace{55\dots 5}_{m_1} \underbrace{22\dots 2}_{m_2} \underbrace{77\dots 7}_{m_3} \underbrace{44\dots 4}_{m_4} \underbrace{11\dots 1}_{m_5} \underbrace{66\dots 6}_{m_6} \underbrace{33\dots 3}_{m_7})$ is the sixth vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_6) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_6 + 3p_0 + 8p_1 + 15p_2 + 8p_3 + 3p_4 - q_4 + 15q_0 + 8q_1 + 3q_2 + 3q_6 + 8q_7 - r_2 + 3r_0 + 3r_4 + 8r_5 + 15r_6 + 8r_7 - s_0 + 3s_2 + 8s_3 + 15s_4 + 8s_5 + 3s_6 - t_6 + 3t_0 + 8t_1 + 15t_2 + 8t_3 + 3t_4 - u_4 + 15u_0 + 8u_1 + 3u_2 + 3u_6 + 8u_7 - w_2 + 3w_0 + 3w_4 + 8w_5 + 15w_6 + 8w_7$, where $\mathbf{c}_6 = (\underbrace{66\dots 6}_{m_1} \underbrace{44\dots 4}_{m_2} \underbrace{22\dots 2}_{m_3} \underbrace{00\dots 0}_{m_4} \underbrace{66\dots 6}_{m_5} \underbrace{44\dots 4}_{m_6} \underbrace{22\dots 2}_{m_7})$ is the seventh vector of $BRep^{m_1+m_2+\dots+m_7}$.

$d_E(\mathbf{x}, \mathbf{c}_7) = m_1 + m_2 + m_3 + m_4 + m_5 + m_6 + m_7 - p_7 + 3p_1 + 8p_2 + 15p_3 + 8p_4 + 3p_5 - q_6 + 3q_0 + 8q_1 + 15q_2 + 8q_3 + 3q_4 - r_5 + 8r_0 + 15r_1 + 8r_2 + 3r_3 + 3r_7 - s_4 + 15s_0 + 8s_1 + 3s_2 + 3s_6 + 8s_7 - t_3 + 8t_0 + 3t_1 + 3t_5 + 8t_6 + 15t_7 - u_2 + 3u_0 + 3u_4 + 8u_5 + 15u_6 + 8u_7 - w_1 + 3w_3 + 8w_4 + 15w_5 + 8w_6 + 3w_7$, where $\mathbf{c}_7 = (\underbrace{77\dots 7}_{m_1} \underbrace{66\dots 6}_{m_2} \underbrace{55\dots 5}_{m_3} \underbrace{44\dots 4}_{m_4} \underbrace{33\dots 3}_{m_5} \underbrace{22\dots 2}_{m_6} \underbrace{11\dots 1}_{m_7})$ is the eighth vector of $BRep^{m_1+m_2+\dots+m_7}$. Thus

$$d(\mathbf{x}, BRep^{m_1+m_2+\dots+m_7}) \leq \frac{11}{2}(m_1 + m_3 + m_5 + m_7) + 6(m_2 + m_6) + 8m_4.$$

Hence the equality.

Theorem 7. $\min\{2m_1+2m_2+2m_3+2m_4+2m_5+2m_6+2m_7, 2m_2+2m_3+2m_4+4m_5+2m_6+2m_7, 2m_1+2m_2+2m_4+2m_5+2m_6+4m_7, 4m_1+2m_2+2m_3+2m_4+2m_6+2m_7, 2m_1+2m_2+4m_3+2m_4+2m_5+2m_6\} \leq r_{HW}(BRep^{m_1+m_2+\dots+m_7}) \leq 11(m_1 + m_3 + m_5 + m_7) + 12(m_2 + m_6) + 16m_4$.

Proof. By choosing $\mathbf{x} = (\underbrace{11\dots\dots\dots 1}_{m_1+m_2+\dots+m_7}) \in \mathbb{Z}_8^{m_1+m_2+\dots+m_7}$ and computing the

homogenous distance from each codeword we get $d_{HW}(\mathbf{x}, BRep^{m_1+m_2+\dots+m_7}) = \min\{2m_1 + 2m_2 + 2m_3 + 2m_4 + 2m_5 + 2m_6 + 2m_7, 2m_2 + 2m_3 + 2m_4 + 4m_5 + 2m_6 + 2m_7, 2m_1 + 2m_2 + 2m_4 + 2m_5 + 2m_6 + 4m_7, 4m_1 + 2m_2 + 2m_3 + 2m_4 + 2m_6 + 2m_7, 2m_1 + 2m_2 + 4m_3 + 2m_4 + 2m_5 + 2m_6\}$. Hence the first inequality follows. The second inequality follows from Proposition 10 and Theorem 6. \square

5 Octonary Simplex Codes of Type α and β

Simplex codes of type α and β have been studied in [26]. The linear code \mathbf{S}_k^α is a type α simplex code over \mathbb{Z}_8 with parameters $(n = 8^k, M = 8^k, d_{HW} = 2^{3(k+1)-2})$ generated by

$$G_k^\alpha = \left[\begin{array}{c|c|c|c|c|c|c|c} 00\dots 0 & 11\dots 1 & 22\dots 2 & 33\dots 3 & 44\dots 4 & 55\dots 5 & 66\dots 6 & 77\dots 7 \\ \hline G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha & G_{k-1}^\alpha \end{array} \right] \quad (21)$$

with $G_1^\alpha = [01234567]$. The number of vectors in \mathbf{S}_k^α is 2^{3k} . The dual code of \mathbf{S}_k^α is denoted by $S_k^{\alpha\perp}$.

The linear code \mathbf{S}_k^β is a type β simplex code over \mathbb{Z}_8 with parameters $(n = 2^{2(k-1)}(2^k - 1), M = 8^k, d_{HW} = 2^{2k-1}(2^k - 1))$ generated by

$$G_2^\beta = \left[\begin{array}{c|c|c|c} 11111111 & 0 & 2 & 4 & 6 \\ \hline 01234567 & 1 & 1 & 1 & 1 \end{array} \right] \quad (22)$$

and for $k > 2$

$$G_k^\beta = \left[\begin{array}{c|c|c|c|c} 11 \cdots 1 & 00 \cdots 0 & 22 \cdots 2 & 44 \cdots 4 & 66 \cdots 6 \\ \hline G_{k-1}^\alpha & G_{k-1}^\beta & G_{k-1}^\beta & G_{k-1}^\beta & G_{k-1}^\beta \end{array} \right], \quad (23)$$

where G_{k-1}^α is the generator matrix of S_{k-1}^α . The dual code of \mathbf{S}_k^β is denoted by $S_k^{\beta\perp}$.

Theorem 8. $r_{HW}(S_k^\alpha) \geq 2^{3k+1}$ and $r_E(S_k^\alpha) \leq 6(8^k - 1) + 2$.

Proof. The proof can be obtained using Proposition 13, Theorem 6, equation (21) and is similar to \mathbb{Z}_4 case [27]. Hence omitted. \square

Theorem 9. $r_E(S_k^\beta) \leq \frac{3}{2}(8^k - 1) - \frac{5}{3}(4^k - 1) - \frac{39}{2} + r_E(S_2^\beta)$ and $r_{HW}(S_k^\beta) \leq 3(8^k - 1) - \frac{10}{3}(4^k - 1) - 139 + r_{HW}(S_2^\beta)$.

Proof. First inequality is proved using Theorem 6 and is similar to \mathbb{Z}_4 case [27]. The case of homogeneous weight is similar. \square

Theorem 10. $r_E(S_k^{\alpha\perp}) \leq 3$, $r_{HW}(S_k^{\alpha\perp}) = 1$ and $r_{HW}(S_k^{\beta\perp}) = 2$.

Proof. By Lemma 4.2 and Theorem 4.3 of [26], $r_E(S_k^{\alpha\perp}) \leq 3$. By Theorem 4.3(3) of [26], $r_{HW}(S_k^{\alpha\perp}) \leq 1$. Since $r_{HW}(S_k^{\alpha\perp}) \geq 1$, so $r_{HW}(S_k^{\alpha\perp}) = 1$. By Theorem 4.4 of [26] and by Theorem 1, $r_{HW}(S_k^{\beta\perp}) \leq 2$ and as $r_{HW}(S_k^{\beta\perp}) \geq 1$ thus $r_{HW}(S_k^{\beta\perp}) = 1$ or 2 but $r_{HW}(S_k^{\beta\perp}) \neq 1$ by Proposition 12. Hence the result follows. \square

Theorem 11. S_k^α and S_k^β are self orthogonal codes over \mathbb{Z}_8 .

Proof. Proof follows from Proposition 3.

6 Octonary MacDonald Codes of Type α and β

The q -ary MacDonald code $\mathbb{M}_{k,u}(q)$ over the finite field \mathbb{F}_q is a unique $[\frac{q^k - q^u}{q - 1}, k, q^{k-1} - q^{u-1}]$ code in which every nonzero codeword has weight either q^{k-1} or $q^{k-1} - q^{u-1}$ [16]. In [12], authors have defined the MacDonald codes over \mathbb{Z}_4 using the generator matrices of simplex codes. In a similar manner one can define MacDonald code over \mathbb{Z}_{2^s} . For $1 \leq u \leq k - 1$, let $G_{k,u}^\alpha$ ($G_{k,u}^\beta$) be the matrix

obtained from $G_k^\alpha \left(G_k^\beta \right)$ by deleting columns corresponding to the columns of $G_u^\alpha \left(G_u^\beta \right)$. i.e,

$$G_{k,u}^\alpha = \left[G_k^\alpha \setminus \frac{\mathbf{0}}{G_u^\alpha} \right], \quad (24)$$

and

$$G_{k,u}^\beta = \left[G_k^\beta \setminus \frac{\mathbf{0}}{G_u^\beta} \right], \quad (25)$$

where $[A \setminus \frac{\mathbf{0}}{B}]$ is the matrix obtained by deleting the matrix $\mathbf{0}$ and B from A where B is a $(k-u) \times 2^{su}$ matrix in (24) and (resp. $(k-u) \times 2^{(s-1)(u-1)}(2^u-1)$) matrix in (25) . The code

$$\mathbb{M}_{k,u}^\alpha : [2^{sk} - 2^{su}, sk] \left(\mathbb{M}_{k,u}^\beta : [2^{(s-1)(u-1)}(2^k-1) - 2^{(s-1)(u-1)}(2^u-1), sk] \right)$$

generated by the matrix $G_{k,u}^\alpha \left(G_{k,u}^\beta \right)$ is the punctured code of $S_k^\alpha \left(S_k^\beta \right)$ and is called a *MacDonald code* of type α (β).

Next theorem provides basic bounds on the covering radii of MacDonald codes over \mathbb{Z}_8 .

Theorem 12.

$$r_E(\mathbb{M}_{k,u}^\alpha) \leq 6(8^k - 8^r) + r_E(\mathbb{M}_{r,u}^\alpha) \text{ for } u < r \leq k,$$

Proof. Similar to \mathbb{Z}_4 case [27].

□

7 Octonary Reed-Muller Code

In this section we give covering radius of octonary first order Reed Muller code [26]. Let $1 \leq i \leq m-2$. Let \mathbf{v}_i be a vector of length 2^{m-2} consisting of successive blocks of 0's and 1's each of size $2^{(m-2)-i}$ and let $\mathbf{1} = (111 \dots 11) \in \mathbf{Z}_2^{2^{m-2}}$. Let G be a $(m-1) \times 2^{m-2}$ matrix given by (consisting of the rows as $\mathbf{1}$ and $4\mathbf{v}_i$ ($1 \leq i \leq m-2$))

$$G = \begin{bmatrix} 0 & 0 & \dots & 0 & 0 & 4 & 4 & \dots & 4 & 4 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 4 & \dots & 0 & 4 & 0 & 4 & \dots & 0 & 4 \\ 1 & 1 & \dots & 1 & 1 & 1 & 1 & \dots & 1 & 1 \end{bmatrix}. \quad (26)$$

The code generated by G is called the *first order Reed-Muller code over \mathbf{Z}_8* , denoted $\mathbb{R}^{1,m-2}$. It is a $(n = 2^{m-2}, M = 2^{m+1}, d_{HW} = 2^{m-1})$ type α linear code over \mathbf{Z}_8 [26]. From Proposition 11 and [31] we have

Theorem 13. $r_{HW}(\mathbb{R}^{1,m-2}) = 2^{m-1} - 2^{\frac{m}{2}-1}$ for even m .

8 Octonary Octa Codes

The octa code over \mathbb{Z}_8 is generated by the following matrix.

$$\mathbb{G} = \begin{bmatrix} 5 & 7 & 5 & 6 & 1 & 0 & 0 & 0 \\ 5 & 0 & 7 & 5 & 6 & 1 & 0 & 0 \\ 5 & 0 & 0 & 7 & 5 & 6 & 1 & 0 \\ 5 & 0 & 0 & 0 & 7 & 5 & 6 & 1 \end{bmatrix}$$

By Proposition 12 we get the following result.

Theorem 14. *If \mathcal{C} is the code generated by G then $r_{HW}(\mathcal{C}) \geq 6$.*

9 Conclusion

In this work, we have introduced new torsion and reduction codes for any linear octonary code and obtained a nice relationship among various reduction and torsion codes. Further we have extended some of the results regarding covering radius of [27] to the octonary case. In particular, we have found exact values and bounds of the covering radius of Repetition codes, Simplex codes of Type α and Type β and their duals, MacDonald codes, and first order Reed-Muller codes, Octacode over \mathbb{Z}_8 . New Reduction and torsion codes can be used to classify octonary linear codes.

References

1. Aoki, T., Gaborit, P., Harada, M., Ozeki, M., and Solé, P.: “On the covering radius of \mathbb{Z}_4 codes and their lattices.” IEEE Trans. Inform. Theory, **vol. 45**, no. 6, 2162–2168, 1999.
2. Arzaki, M., and Suprijanto, D. *Note on Elementary Constructions of Self-Dual Codes over \mathbb{Z}_8* . International Mathematical Forum, **Vol. 6, No. 16**, 785-794, (2011).
3. Bannai E., Dougherty S.T., Harada M. and Oura M.: *Type II codes, even unimodular lattices and invariant rings*. IEEE Trans. Inform. Theory, **45**, 1194–1205, (1999).
4. Berlekamp, E.R., McEliece, R.J., van Tilborg, H.C.A. *On the Inherent Intractability of some Coding Problems*. IEEE Trans. Inform. Theory, **24(3)**, 384-386, (1996).
5. Betsumiya, K., Betty, R. A. L., Munemasa, A. *Mass Formula for Even Codes over \mathbb{Z}_8* . Cryptography and Coding, M.G. Parker(Ed.), LNCS 5921, 65-77, (2009).
6. Bonnetcaze A., Solé P. and Calderbank A.R., *Quaternary quadratic residue codes and unimodular lattices*. IEEE Trans. Inform. Theory **41**, 366–377, (1995).
7. Bonnetcaze A., Solé P., Bachoc C. and Mourrain B., *Type II codes over \mathbb{Z}_4* . IEEE Trans. Inform. Theory **43**, 969–976, (1997).
8. C. Carlet, “ \mathbb{Z}_{2^k} -linear codes.” IEEE Trans. Inform. Theory, **vol. 44**, no. 4, 1543–1547, (1998).
9. Chiu, M.H., Yau, S.S.T., and Yu, Y.: *\mathbb{Z}_8 -Cyclic Codes and Quadratic Residue Codes*. Advances in Applied Mathematics, **25**, 12-33, (2000).
10. Cohen, G. D., Karpovsky, M. G., Mattson, H. F. and Schatz, J. R.: “Covering radius-Survey and recent results.” IEEE Trans. Inform. Theory, **vol. 31**, no. 3, 328–343, (1985).

11. Cohen C., Honkala I., Litsyn, S., and Lobstein A., **Covering Codes**, Elsevier, 1997.
12. Colbourn, C.J. and Gupta, M.K.: “ *On quaternary MacDonal codes.*” Proc. Information Technology: Coding and Computing (ITCC), April 2003, 212–215.
13. I. Constantinescu, W. Heise, and T. Honold “ *Monomial extensions of isometries between codes over \mathbb{Z}_m .*” Proc. Workshop ACCT’96, Sozopol, Bulgaria, 1996, 98–104.
14. Conway J.H. and Sloane N.J.A., *Self-dual codes over the integers modulo 4.* J. Combin. Theory Ser. A **62**, 30–45, (1993).
15. J. A. Davis and J. Jedwab, “*Peak to mean power control in OFDM, Golay complementary sequences, and Reed-Muller codes.*” IEEE Trans. Inform. Theory, **vol. 45**, no. 7, 2397–2417, (1999).
16. S. Dodunekov and J. Simonis. *Codes and projective multisets.* The Electronic Journal of Combinatorics **5** (1998) R37.
17. Dougherty S.T., Gulliver T.A. and Harada M., *Type II codes over finite rings and even unimodular lattices.* J. Alg. Combin., **9**, 233–250, (1999).
18. Dougherty S.T., Harada M. and Solé P., *Self-dual codes over rings and the Chinese Remainder Theorem.* Hokkaido Math. J., **28**, 253–283, (1999).
19. Dougherty S.T., Harada M. and Solé P., *Shadow codes over \mathbb{Z}_4 .* Finite Fields and Their Appl., **7**, 507–529, (2001).
20. Dougherty S.T., Gulliver, T.A., and Wong, A.: *Self-dual codes over \mathbb{Z}_8 and \mathbb{Z}_9 ,* Design Codes and Cryptography. **41**, 235-249, (2006).
21. Durairajan, C. **On Covering Codes and Covering radius of some optimal codes**, PhD Thesis, Department of Mathematics, IIT Kanpur, (1996).
22. Garg, A., and Dutt, S. *Cyclic Codes of length 2^k over \mathbb{Z}_8 .* Open Journal of Applied Sciences, 104-107, (2012).
23. Gulliver T.A. and Harada M., *Double circulant self dual codes over \mathbb{Z}_{2k} .* IEEE Trans. Inform. Theory, **44**, 3105–3123, (1998).
24. Gulliver, T.A., and Harada, M. *Extremal Self-Dual Codes over \mathbb{Z}_6 , \mathbb{Z}_8 , and \mathbb{Z}_{10} .* AKCE J. Graphs. Combin., **2**, No. 1, 11-24, (2005).
25. Glynn, D., Gulliver, T.A. and Gupta, M.K. *On Some Quaternary Self-Orthogonal Codes.* Ars. Comb., **Vol 85**, 193-209, (2007).
26. Gupta, M.K., *On Some Linear Codes over \mathbb{Z}_{2^s} ,* PhD Thesis, Department of Mathematics, IIT Kanpur, (1999).
27. Gupta, M.K. and Durairajan C., *On the covering radius of some modular codes.* Advances in Mathematics of Communications, **Vol 8**, No. 2, 129-137, (2014).
28. Hammons, A. R., Kumar, P. V., Calderbank, A. R., Sloane, N. J. A. and Solé, P.: *The \mathbb{Z}_4 -linearity of kerdock, preparata, goethals, and related codes.* IEEE Trans. Inform. Theory, **40**, 301–319, (1994).
29. Hammons, Jr. A.R., Kumar P.V., Calderbank A.R., Sloane N.J.A. and Solé P., *A linear construction for certain Kerdock and Preparata codes.* Bull Amer. Math. Soc. **29**, 218–222, (1993).
30. Harada M., *New extremal Type II codes over \mathbb{Z}_4 .* Des. Codes and Cryptogr. **13**, 271–284, (1998).
31. Helleseth, T., Klove, T. and Mykkelveit, J.: *On the Covering Radius of Binary Codes,* IEEE Trans. Inform. Theory, **Vol. 24**, 627-628, (1978).
32. Piret, P.M. *Algebraic Construction of Cyclic Codes over \mathbb{Z}_8 with a Good Euclidean Minimum Distance.* IEEE Trans. on Inform. Theory, **Vol 41**, No. 3, 815-818, (1995).
33. Rains, E. M. and Sloane, N. J. A. *Self-Dual Codes* in V. Pless and W.C. Huffman (Eds.) **The Handbook of Coding Theory**, North-Holland, New York, (1998).

34. Yildiz, B., Siap, I., Bilgin, T., and Yesilot, G. *The covering problem for finite rings with respect to the RT -metric*. Applied Mathematics Letters, **Vol 23, Issue 9**, 988-992, (2010).